

MATH NEWS



Grade 5, Module 3, Topic C

5th Grade Math

Module 3: Addition and Subtraction of Fractions

Math Parent Letter

This document is created to give parents and students an understanding of the math concepts found in Eureka Math (© 2013 Common Core, Inc.) that is also posted as the Engage New York material which is taught in the classroom. Grade 5 Module 3 of Eureka Math (Engage New York) covers Addition and Subtraction of Fractions. This newsletter will discuss Module 3, Topic C.

Topic C: Making Like Units Numerically

Words to know:

- equivalence
- numericallysum
- difference

mixed number

- - improper fraction

Things to Remember!!!

- Equivalence being equal, having the same value
- Numerically using numbers
- Sum the answer to an addition problem
- Difference the answer to a subtraction problem
- **Number Line** a line used to show placement of whole numbers, fractions, and mixed numbers
- Mixed Number a whole number plus a fraction smaller than 1, written without the + sign, e.g. $5\frac{3}{4}$ means $5 + \frac{3}{4}$
- Improper Fraction a fraction with the numerator equal to or greater than the denominator


OBJECTIVES OF TOPIC C


- Add fractions to and subtract fractions from **whole numbers** using **equivalence** and the **number line** as strategies.
- Add fractions making like units **numerically.**
- Add fractions with **sums** greater than 2.
- Subtract fractions making like units numerically.
- Subtract fractions greater than or equal to 1.

Focus Area- Topic C: Making Like Units


Problem 1: 2 + $2\frac{1}{2} = 4\frac{1}{2}$

Step 1: Add the whole numbers. **Step 2**: Add the fraction.





Step 1: Subtract the whole numbers. **Step 2**: Subtract the fraction.



Problem 3: $3 = \frac{3}{4} - \frac{1}{5} = \left(\frac{3x5}{4x5}\right) + \left(\frac{1x4}{5x4}\right)$ Step 1: Make like units $\frac{15}{20} + \frac{4}{20} = \frac{19}{20}$ numerically.

Step 2: Add fractions.

Problem 4:	\rightarrow $7\frac{5}{8} + 8\frac{2}{5}$
Step 1 : Add the whole numbers.	$=7+8+\frac{5}{8}+\frac{2}{5}$
Step 2: Make like units numerically.	$= 15 + \left(\frac{5 \times 5}{8 \times 5}\right) + \left(\frac{2 \times 8}{5 \times 8}\right)$
Step 3: Add fractions.	$= 15 + \frac{25}{40} + \frac{16}{40}$
Step 4 : If sum is an improper fraction, rename fraction as a mixed	$= 15 + \frac{41}{40}$
number. Step 5 : Add whole	$= 15 + 1 + \frac{1}{40}$
number to fraction.	$= 16 \frac{1}{40}$
Step 6 : Simplify sum if possible.	10

Problem 5:	$5\frac{2}{3} - 2\frac{1}{2}$	
	$= (5-2) + \frac{2}{3} - \frac{1}{2}$	(Step 1: Subtract the whole numbers.)
	$= 3 + \frac{2}{3} - \frac{1}{2}$	
	$= (3-\frac{1}{2})+\frac{2}{3}$	(Step 2: Subtract the second fraction from the whole number.)
	$= 2\frac{1}{2} + \frac{2}{3}$	(Step 3: Make like units numerically.)
	$= 2 + \left(\frac{1x3}{2x3}\right) + \left(\frac{2x2}{3x2}\right)$	
	$= 2 + \frac{3}{6} + \frac{4}{6}$	(Step 4: Add the fractions.)
	$= 2 + \frac{7}{6}$	(Step 5: If sum of the fractions is an improper fraction, rename as a whole or mixed number.)
	$=$ 2 + 1 + $\frac{1}{6}$	(Step 6: Add fraction to whole numbers.)
	$= 3\frac{1}{6}$	(Step 7: Simplify fraction if possible.)

Problem 6: Mrs. Sanchez made $7\frac{4}{5}$ gallons of punch for a party. If there were $10\frac{1}{2}$ gallons in the mixture, how many gallons did she have left in the mixture?

$$10\frac{1}{2} - 7\frac{4}{5}$$

$$= (10 - 7) + \frac{1}{2} - \frac{4}{5}$$

$$= 3 + \frac{1}{2} - \frac{4}{5}$$

$$= (3 - \frac{4}{5}) + \frac{1}{2}$$

$$= 2\frac{1}{5} + \frac{1}{2}$$

$$= 2 + (\frac{1x^2}{5x^2}) + (\frac{1x^5}{2x^5})$$

$$= 2 + \frac{2}{10} + \frac{5}{10} = 2\frac{7}{10}$$
There are $2\frac{7}{10}$ gallons of Mrs. Sanchez's punch mixture left.

Problem 7: Bryant has a goal to drink at least $6\frac{1}{2}$ quarts of water during his day of training for the big marathon race. On his first break he drank $1\frac{3}{4}$ quarts, and during his second break he had another $2\frac{1}{5}$ quarts. How many quarts of water should Bryant drink on his last break of the day to reach his goal?

$$6\frac{1}{2} - \left(1\frac{3}{4} + 2\frac{1}{5}\right) = 6\frac{1}{2} - \left(3\frac{3}{4} + \frac{1}{5}\right) = 6\frac{1}{2} - \left(3 + \frac{3x5}{4x5} + \frac{1x4}{5x4}\right) = 6\frac{1}{2} - \left(3 + \frac{15}{20} + \frac{4}{20}\right)$$

$$6\frac{1}{2} - 3\frac{19}{20} = (6 - 3) + \frac{1}{2} - \frac{19}{20} = 3 + \frac{1}{2} - \frac{19}{20} = (3 - \frac{19}{20}) + \frac{1}{2}$$

$$2\frac{1}{20} + \frac{1}{2} = 2 + \left(\frac{1x2}{20x2}\right) + \left(\frac{1x20}{2x20}\right) = 2 + \frac{2}{40} + \frac{20}{40} = 2\frac{22}{40} = 2\frac{22+2}{40+2} = 2\frac{11}{20}$$
Or
$$2\frac{1}{20} + \frac{1}{2} = 2 + \left(\frac{1x1}{20x1}\right) + \left(\frac{1x10}{2x10}\right) = 2 + \frac{1}{20} + \frac{10}{20} = 2\frac{11}{20}$$
Bryant should drink $2\frac{11}{20}$ quarts of water to reach his goal.

Students do **not** have to use the least common denominator. They are just expected to create common denominators. In the end the answers will be the same.

**** The strategy above is a possible approach. The student could have first added $1\frac{3}{4} + 2\frac{1}{5}$. Then take the sum and subtract from $6\frac{1}{2}$.